



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

REC'D 25 AUG 2003

WIPO

PCT

PCT/EP 03/07173 Rec'd PCT/PTC 05 JAN 2005 #2

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02015063.7

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 02015063.7
Demande no:

Anmeldetag:
Date of filing: 05.07.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

EVOTEC Neurosciences GmbH
Schnackenburgallee 114
22525 Hamburg
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Diagnostic and therapeutic use of TB2 gene and protein for neurodegenerative diseases

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

G01N33/48

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

DIAGNOSTIC AND THERAPEUTIC USE OF TB2 GENE AND PROTEIN FOR NEURODEGENERATIVE DISEASES

The present invention relates to methods of diagnosing, prognosticating, and monitoring the progression of neurodegenerative diseases in a subject. Furthermore, methods of therapy control and screening for modulating agents of neurodegenerative diseases are provided. The invention also discloses pharmaceutical compositions, kits, and recombinant animal models.

Neurodegenerative diseases, in particular Alzheimer's disease (AD), have a strongly debilitating impact on a patient's life. Furthermore, these diseases constitute an enormous health, social and economic burden. AD is the most common age-related neurodegenerative condition affecting about 10 % of the population over 65 years of age and up to 45 % over age 85 (for a recent review see Vickers et al., *Progress in Neurobiology* 2000, 60: 139-165). Presently this amounts to an estimated 12 million cases in the US, Europe, and Japan. This situation will inevitably worsen with the demographic increase in the number of old people ("aging of the baby boomers") in developed countries. The neuropathological hallmarks that occur in the brains of individuals with AD are senile plaques, composed of amyloid- β protein, and profound cytoskeletal changes coinciding with the appearance of abnormal filamentous structures and the formation of neurofibrillary tangles. AD is a progressive disease that is associated with early deficits in memory formation and ultimately leads to the complete erosion of higher cognitive function. A characteristic feature of the pathogenesis of AD is the selective vulnerability of particular brain regions and subpopulations of nerve cells to the degenerative process. Specifically, the temporal lobe region and the hippocampus are affected early and more severely during the progression of the disease. On the other hand, neurons within the frontal cortex, occipital cortex, and the cerebellum remain largely intact and are protected from neurodegeneration (Terry et al., *Annals of Neurology* 1981, 10: 184-192).

Currently, there is no cure for AD, nor is there an effective treatment to halt the progression of AD or even to diagnose AD ante-mortem with high probability. Several risk factors have been identified that predispose an individual to develop AD, among them most prominently the epsilon4 allele of apolipoprotein E (ApoE). Although there are rare examples of early-onset AD which have been attributed to genetic defects in the genes for amyloid precursor protein (APP), presenilin-1, and presenilin-2, the prevalent form of late-onset sporadic AD is of hitherto unknown etiologic origin. The late onset and complex pathogenesis of neurodegenerative disorders pose a formidable challenge to the development of therapeutic and diagnostic agents. It is crucial to expand the pool of potential drug targets and diagnostic markers. It is therefore an object of the present invention to provide insight into the pathogenesis of neurological diseases and to provide methods, materials, and animal models which are suited inter alia for the diagnosis and development of a treatment of these diseases. This object has been solved by the features of the independent claims. The subclaims define preferred embodiments of the present invention.

The present invention discloses the differential expression of the human gene coding for TB2 in Alzheimer's disease brain samples by using suppressive subtractive hybridization and screening of DNA biochips. The human gene encoding TB2 (alias: polyposis locus protein 1, DP1, deleted in polyposis 1) was first described in attempts to dissect the familial adenomatous polyposis locus on chromosome 5q21. Familial adenomatous polyposis (FAP) is an autosomal-dominant, inherited disease in which affected individuals develop adenomatous polyps within their colon. Some of these polyps may eventually progress to malignancy, thus constituting a life-threatening disease condition. It is thought that the FAP locus on chromosome 5q21 contains tumor suppressor genes that, when deleted or mutated, fail to control cell proliferation and cause neoplasia. The FAP locus was found to contain at least six genes termed FER, MCC, SRP19, TB1, TB2, and APC (Kinzler et al., *Science* 1991, 253: 661-665). A second study found three genes in a 100 kb deleted segment on chromosome 5 in two unrelated patients with FAP (Joslyn

et al., *Cell* 1991, 66: 601-613). These three genes were named SRP19, DP1, and APC, wherein DP1 is identical to TB2 of the aforementioned study. Mutations in the APC gene were detected in FAP patients as well as in sporadic colorectal cancer patients. Consequently, APC is regarded as the candidate tumor suppressor gene within the FAP locus (US 5352775 / US Re. 36713; US 5783666; US 5648212; US 6114124).

Human TB2 is a putative membrane protein (nucleotide sequence of mRNA: NCBI GenBank accession number BC000232; amino acid sequence of TB2 protein: NCBI GenBank accession numbers Q00765, AAH00232) comprising 185 amino acids with at least two predicted transmembrane domains. It is a member of an evolutionary conserved family of proteins including a mouse ortholog with 92 % identity at the amino acid level (Prieschl et al., *Gene* 1996, 169: 215-218), a yeast homolog named Yop1p (Calero et al., *J. Biol. Chem.* 2001, 276: 12100-12112) and further homologs from *C. elegans*, *D. melanogaster*, and *P. falciparum*. The stress-inducible plant homolog HVA22 falls into the same protein family (Shen et al., *Plant Mol. Biol.* 2001, 45: 327-340). The function of human TB2 is largely unknown but may be inferred from studies conducted on the *S. cerevisiae* Yop1p protein. Yop1p was identified as an interacting protein for Yip1p, which plays a role in Rab GTPase mediated membrane transport in yeast (Calero et al., *J. Biol. Chem.* 2001, 276: 12100-12112). Yop1p is a Golgi membrane protein with a hydrophilic region at its N-terminus through which it interacts specifically with the cytosolic domain of Yip1p. Yop1p was also copurified with a number of Rab GTPases from cellular lysates. Yop1p overexpression profoundly compromised yeast cell growth and led to an accumulation of internal cell membranes and a block in membrane traffic. Therefore, it was followed that Yop1p together with Yip1p plays an important role in cellular membrane traffic (Calero et al., *J. Biol. Chem.* 2001, 276: 12100-12112).

The tissue expression of human TB2 is broad. Its mRNA has been found in a variety of tissues including colonic mucosa, skin/melanocytes, placenta, thyroid gland, epithelial cells of the lung, the cochlea of the ear, and the amygdala (Source: GeneReport number D5S346). The murine ortholog is expressed ubiquitously (Prieschl et al., *Gene* 1996, 169: 215-218). The latter

study reported a downregulation of TB2 mRNA upon stimulation of mast cells by IgE plus antigen (ibid.). An independent study detected a prominent level of TB2 mRNA in mouse brain but not in liver (Kawai et al., *Biochem. Biophys. Res. Comm.* 1995, 213: 967-974).

Based on the above information it is reasonable to assume that TB2, as an integral membrane protein, participates in cellular membrane traffic involving the Golgi apparatus. It may exert its function in many tissues including the amygdala of the brain. TB2 may, for instance, impact on sorting and processing of the amyloid precursor protein and, thereby, have a causative role in the development of AD.

The present invention discloses a dysregulation of Tb2 gene expression on the transcriptional level in the temporal cortex region relative to the frontal lobe region of brain samples taken from AD patients. No such dysregulation is observed in samples derived from age-matched, healthy controls. To date, no experiments have been described that demonstrate a relationship between a dysregulation of TB2 gene expression and the pathology of neuro-degenerative diseases, in particular AD. Such a link, as disclosed in the present invention, offers new ways, inter alia, for the diagnosis and treatment of said diseases.

The singular forms "a", "an", and "the" as used herein and in the claims include plural reference unless the context dictates otherwise. For example, "a cell" means as well a plurality of cells, and so forth. The term "and/or" as used in the present specification and in the claims implies that the phrases before and after this term are to be considered either as alternatives or in combination. For instance, the wording "determination of a level and/or an activity" means that either only a level, or only an activity, or both a level and an activity are determined. The term "level" as used herein is meant to comprise a gage of, or a measure of the amount of, or a concentration of a transcription product, for instance an mRNA, or a translation product, for instance a protein or polypeptide. The term "activity" as used herein shall be understood as a measure for the ability of a transcription product or a

translation product to produce a biological effect or a measure for a level of biologically active molecules. The term "activity" also refers to enzymatic activity. The terms "level" and/or "activity" as used herein further refer to gene expression levels or gene activity. Gene expression can be defined as the utilization of the information contained in a gene by transcription and translation leading to the production of a gene product. A gene product comprises either RNA or protein and is the result of expression of a gene. The amount of a gene product can be used to measure how active a gene is. The term "gene" as used in the present specification and in the claims comprises both coding regions (exons) as well as non-coding regions (e.g. non-coding regulatory elements such as promoters or enhancers, introns, leader and trailer sequences). "Regulatory elements" shall comprise inducible and non-inducible promoters, enhancers, operators, and other elements that drive and regulate gene expression. The term "fragment" as used herein is meant to comprise e.g. an alternatively spliced, or truncated, or otherwise cleaved transcription product or translation product. The term "derivative" as used herein refers to a mutant, or an RNA-edited, or a chemically modified, or otherwise altered transcription product, or to a mutant, or chemically modified, or otherwise altered translation product. For instance, a "derivative" may be generated by processes such as altered phosphorylation, or glycosylation, or, acetylation, or lipidation, or by altered signal peptide cleavage or other types of maturation cleavage. These processes may occur post-translationally. The term "modulator" as used in the present invention and in the claims refers to a molecule capable of changing or altering the level and/or the activity of a gene, or a transcription product of a gene, or a translation product of a gene. Preferably, a "modulator" is capable of changing or altering the biological activity of a transcription product or a translation product of a gene. Said modulation, for instance, may be an increase or a decrease in enzyme activity, a change in binding characteristics, or any other change or alteration in the biological, functional, or immunological properties of said translation product of a gene. The terms "agent", "reagent", or "compound" refer to any substance, chemical, composition or extract that have a positive or negative biological effect on a cell, tissue, body fluid, or within the context of any

biological system, or any assay system examined. They can be agonists, antagonists, partial agonists or inverse agonists of a target. Such agents, reagents, or compounds may be nucleic acids, natural or synthetic peptides or protein complexes, or fusion proteins. They may also be antibodies, organic or anorganic molecules or compositions, small molecules, drugs and any combinations of any of said agents above. They may be used for testing, for diagnostic or for therapeutic purposes. The terms "oligonucleotide primer" or "primer" refer to short nucleic acid sequences which can anneal to a given target polynucleotide by hybridization of the complementary base pairs and can be extended by a polymerase. They may be chosen to be specific to a particular sequence or they may be randomly selected, e.g. they will prime all possible sequences in a mix. The length of primers used herein may vary from 10 nucleotides to 80 nucleotides. "Probes" are short nucleic acid sequences of the nucleic acid sequences described and disclosed herein or sequences complementary therewith. They may comprise full length sequences, or fragments, derivatives, isoforms, or variants of a given sequence. The identification of hybridization complexes between a "probe" and an assayed sample allows the detection of the presence of other similar sequences within that sample. The term "variant" as used herein refers to any polypeptide or protein, in reference to polypeptides and proteins disclosed in the present invention, in which one or more amino acids are added and/or substituted and/or deleted and/or inserted at the N-terminus, and/or the C-terminus, and/or within the native amino acid sequences of the native polypeptides or proteins of the present invention. Furthermore, the term "variant" shall include any shorter or longer version of a polypeptide or protein. "Variants" shall also comprise a sequence that has at least about 80% sequence identity, more preferably at least about 90% sequence identity, and most preferably at least about 95% sequence identity with the amino acid sequences of translation products of a gene coding for Tb2. "Variants" of such a protein molecule include, for example, proteins with conservative amino acid substitutions in highly conservative regions. "Proteins and polypeptides" of the present invention include variants, fragments and chemical derivatives of the Tb2 protein. They can include proteins and polypeptides which can be

isolated from nature or be produced by recombinant and/or synthetic means. Native proteins or polypeptides refer to naturally-occurring truncated or secreted forms, naturally occurring variant forms (e.g. splice-variants) and naturally occurring allelic variants. In the present invention, the terms "risk", "susceptibility", and "predisposition" are tantamount and are used with respect to the probability of developing a neurodegenerative disease, preferably Alzheimer's disease. The term 'AD' shall mean Alzheimer's disease.

Neurodegenerative diseases or disorders according to the present invention comprise Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, Pick's disease, fronto-temporal dementia, progressive nuclear palsy, corticobasal degeneration, cerebro-vascular dementia, multiple system atrophy, and mild-cognitive impairment. Further conditions involving neurodegenerative processes are, for instance, ischemic stroke, age-related macular degeneration, narcolepsy, motor neuron diseases, nerve injury and repair, and multiple sclerosis.

In one aspect, the invention features a method of diagnosing or prognosticating a neurodegenerative disease in a subject, or determining whether a subject is at increased risk of developing said disease. The method comprises: determining a level, or an activity, or both said level and said activity of (i) a transcription product of the TB2 gene, and/or of (ii) a translation product of the TB2 gene, and/or of (iii) a fragment or derivative of said transcription or translation product in a sample from said subject and comparing said level, and/or said activity to a reference value representing a known disease or health status, thereby diagnosing or prognosticating said neurodegenerative disease in said subject, or determining whether said subject is at increased risk of developing said neurodegenerative disease.

In a further aspect, the invention features a method of monitoring the progression of a neurodegenerative disease in a subject. A level, or an activity, or both said level and said activity, of (i) a transcription product of the TB2 gene, and/or of (ii) a translation product of the TB2 gene, and/or of (iii) a

fragment or derivative of said transcription or translation product in a sample from said subject is determined. Said level and/or said activity is compared to a reference value representing a known disease or health status. Thereby the progression of said neurodegenerative disease in said subject is monitored.

In still a further aspect, the invention features a method of evaluating a treatment for a neurodegenerative disease, comprising determining a level, or an activity, or both said level and said activity of (i) a transcription product of the TB2 gene, and/or of (ii) a translation product of the TB2 gene, and/or of (iii) a fragment or derivative of said transcription or translation product in a sample obtained from a subject being treated for said disease. Said level, or said activity, or both said level and said activity are compared to a reference value representing a known disease or health status, thereby evaluating the treatment for said neurodegenerative disease.

In a preferred embodiment, said subjects suffer from Alzheimer's disease.

The present invention discloses the differential expression and regulation of the TB2 gene in specific brain regions of AD patients. Consequently, the TB2 gene and its corresponding transcription and/or translation products may have a causative role in the regional selective neuronal degeneration typically observed in AD. Alternatively, the TB2 protein may confer a neuroprotective function to the remaining surviving nerve cells. Based on these disclosures, the present invention has utility for the diagnostic evaluation and prognosis as well as for the identification of a predisposition to a neurodegenerative disease, in particular AD. Furthermore, the present invention provides methods for the diagnostic monitoring of patients undergoing treatment for such a disease.

It is particularly preferred that said sample to be analyzed and determined is selected from the group consisting of a brain tissue or other tissues, organs or body cells. The sample can also consist of cerebrospinal fluid or other body fluids such as saliva, urine, blood, serum plasma, or nasal mucosa.

In further preferred embodiments, said reference value is that of a level, or an activity, or both said level and said activity of (i) a transcription product of the TB2 gene, and/or of (ii) a translation product of the TB2 gene, and/or of (iii) a fragment or derivative of said transcription or translation product in a sample from a subject not suffering from said neurodegenerative disease.

In preferred embodiments, an alteration in the level and/or activity of a transcription product of a gene coding for TB2 and/or a translation product of a gene coding for TB2 in a sample cell, or tissue, or body fluid from said subject relative to a reference value representing a known health status indicates a diagnosis, or prognosis, or increased risk of becoming diseased with a neurodegenerative disease, particularly AD.

In preferred embodiments, the measurement of the level of transcription product of the TB2 gene is performed in a sample from a subject using a quantitative PCR-analysis with primer combinations to amplify said gene specific sequences from cDNA obtained by reverse transcription of RNA extracted from a sample of a subject. A Northern blot with probes specific for said gene can also be applied. It might further be preferred to measure transcription products by means of chip-based micro-array technologies. These techniques are known to those of ordinary skill in the art (see Sambrook and Russell, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2000).

Furthermore, a level and/or activity of a translation product of the TB2 gene and/or fragment of said translation product, can be detected using an immunoassay, an activity assay and/or binding assay. These assays can measure the amount of binding between said protein molecule and an anti-protein antibody by the use of enzymatic, chromodynamic, radioactive, or luminescent labels which are attached to either the anti-protein antibody or a secondary antibody which binds the anti-protein antibody. In addition, other high affinity ligands may be used. Immunoassays which can be used include

e.g. ELISAs, Western blots and other techniques known to those of ordinary skill in the art (see Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1999). All of these detection techniques may also be employed in the format of micro-arrays, protein-arrays, or protein-chip based technologies.

In a preferred embodiment, the level, or the activity, or both said level and said activity of (i) a transcription product of the TB2 gene, and/or of (ii) a translation product of the TB2 gene, and/or of (iii) a fragment or derivative of said transcription or translation product in a series of samples taken from said subject over a period of time is compared, in order to monitor the progression of said disease. In further preferred embodiments, said subject receives a treatment prior to one or more of said sample gatherings. In yet another preferred embodiment, said level and/or activity is determined before and after said treatment of said subject.

In another aspect, the invention features a kit for diagnosing or prognosticating neurodegenerative diseases, in particular AD, in a subject, or determining the propensity or predisposition of a subject to develop a neurodegenerative disease, in particular AD, said kit comprising:

(a) at least one reagent which is selected from the group consisting of (i) reagents that selectively detect a transcription product of the TB2 gene (ii) reagents that selectively detect a translation product of the TB2 gene; and

(b) an instruction for diagnosing, or prognosticating a neurodegenerative disease, in particular AD, or determining the propensity or predisposition of a subject to develop such a disease by

- detecting a level, or an activity, or both said level and said activity, of said transcription product and/or said translation product of the TB2 gene, in a sample from said subject; and
- diagnosing or prognosticating a neurodegenerative disease, in particular AD, or determining the propensity or predisposition of said subject to develop such a disease,

wherein a varied level, or activity, or both said level and said activity, of said transcription product and/or said translation product compared to a reference value representing a known health status; or a level, or activity, or both said level and said activity, of said transcription product and/or said translation product similar or equal to a reference value representing a known disease status, indicates a diagnosis or prognosis of a neurodegenerative disease, in particular AD, or a increased propensity or predisposition of developing such a disease. The kit, according to the present invention, may be particularly useful for the identification of individuals that are at risk of developing a neurodegenerative disease, in particular AD. Consequently, the kit, according to the invention, may serve as a means for targeting identified individuals for early preventive measures or therapeutic intervention prior to disease onset, before irreversible damage in the course of the disease has been inflicted. Furthermore, in preferred embodiments, the kit featured in the invention is useful for monitoring a progression of a neurodegenerative disease, in particular AD, in a subject, as well as monitoring success or failure of therapeutic treatment for such a disease of said subject.

In another aspect, the invention features a method of treating or preventing a neurodegenerative disease, in particular AD, in a subject comprising the administration to said subject in a therapeutically or prophylactically effective amount of an agent or agents which directly or indirectly affect a level, or an activity, or both said level and said activity, of (i) the TB2 gene, and/or (ii) a transcription product of the TB2 gene, and/or (iii) a translation product of the TB2 gene, and/or (iv) a fragment or derivative of (i) to (iii). Said agent may comprise a small molecule, or it may also comprise a peptide, an oligopeptide, or a polypeptide. Said peptide, oligopeptide or polypeptide may comprise an amino acid sequence of a translation product of a gene coding for TB2, or a fragment, or derivative, or a variant thereof.

In preferred embodiments, the method comprises the application of per se known methods of gene therapy and/or antisense nucleic acid technology to administer said agent or agents. In general, gene therapy includes several

approaches: molecular replacement of a mutated gene, addition of a new gene resulting in the synthesis of a therapeutic protein, and modulation of endogenous cellular gene expression by recombinant expression methods or by drugs. Gene-transfer techniques are described in detail (see e.g. Behr, *Acc Chem Res* 1993, 26: 274-278 and Mulligan, *Science* 1993, 260: 926-93) and include direct gene-transfer techniques such as mechanical microinjection of DNA into a cell as well as indirect techniques employing biological vectors (like recombinant viruses, especially retroviruses) or model liposomes, or techniques based on transfection with DNA coprecipitation with polycations, cell membrane perturbation by chemical (solvents, detergents, polymers, enzymes) or physical means (mechanic, osmotic, thermic, electric shocks). The postnatal gene transfer into the central nervous system has been described in detail (see e.g. Wolff, *Curr Opin Neurobiol* 1993, 3: 743-748).

In particular, the invention features a method of treating or preventing a neurodegenerative disease by means of antisense nucleic acid therapy, i.e. the down-regulation of an inappropriately expressed or defective gene by the introduction of antisense nucleic acids or derivatives thereof into certain critical cells (see e.g. Gillespie, *DN&P* 1992, 5: 389-395; Agrawal and Akhtar, *Trends Biotechnol* 1995, 13: 197-199; Crooke, *Biotechnology* 1992, 10: 882-886). Apart from hybridization strategies, the application of ribozymes, i.e. RNA molecules that act as enzymes, destroying RNA that carries the message of disease has also been described (see e.g. Barinaga, *Science* 1993, 262: 1512-1514). In preferred embodiments, the subject to be treated is a human, and therapeutic antisense nucleic acids or derivatives thereof are directed against transcripts of a gene coding for TB2. It is preferred that cells of the central nervous system, preferably the brain, of a subject are treated in such a way. Cell penetration can be performed by known strategies such as coupling of antisense nucleic acids and derivatives thereof to carrier particles, or the above described techniques. Strategies for administering targeted therapeutic oligodeoxynucleotides are known to those of skill in the art (see e.g. Wickstrom, *Trends Biotechnol* 1992, 10: 281-287). In some cases, delivery can be performed by mere topical application. Further approaches

are directed to intracellular expression of antisense RNA. In this strategy, cells are transformed *ex vivo* with a recombinant gene that directs the synthesis of an RNA that is complementary to a region of target nucleic acid. Therapeutical use of Intracellularly expressed antisense RNA is procedurally similar to gene therapy.

In further preferred embodiments, the method comprises grafting donor cells into the central nervous system, preferably the brain, of said subject, or donor cells preferably treated so as to minimize or reduce graft rejection, wherein said donor cells are genetically modified by insertion of at least one transgene encoding said agent or agents. Said transgene might be carried by a viral vector, in particular a retroviral vector. The transgene can be inserted into the donor cells by a nonviral physical transfection of DNA encoding a transgene, in particular by microinjection. Insertion of the transgene can also be performed by electroporation, chemically mediated transfection, in particular calcium phosphate transfection, liposomal mediated transfection.

In preferred embodiments, said agent for treating or preventing a neurodegenerative disease, in particular AD, is a therapeutic protein which can be administered to said subject, preferably a human, by a process comprising introducing subject cells into said subject, said subject cells having been treated *in vitro* to insert a DNA segment encoding said therapeutic protein, said subject cells expressing *in vivo* in said subject a therapeutically effective amount of said therapeutic protein. Said DNA segment can be inserted into said cells *in vitro* by a viral vector, in particular a retroviral vector.

Methods of treatment, according to the present invention, comprise the application of therapeutic cloning, transplantation, and stem cell therapy using embryonic stem cells or embryonic germ cells and neuronal adult stem cells, combined with any of the previously described cell- and gene therapeutic methods. Stem cells may be totipotent or pluripotent. They may also be organ-specific. Strategies for repairing diseased and/or damaged brain cells

or tissue comprise (i) taking donor cells from an adult tissue. Nuclei of those cells are transplanted into unfertilized egg cells from which the genetic material has been removed. Embryonic stem cells are isolated from the blastocyst stage of the cells which underwent somatic cell nuclear transfer. Use of differentiation factors then leads to a directed development of the stem cells to specialized cell types, preferably neuronal cells (Lanza et al., *Nature Medicine* 1999, 9: 975-977), or (ii) purifying adult stem cells, isolated from the central nervous system, or from bone marrow (mesenchymal stem cells), for in vitro expansion and subsequent grafting and transplantation, or (iii) directly inducing endogenous neural stem cells to proliferate, migrate, and differentiate into functional neurons (Peterson DA, *Curr. Opin. Pharmacol.* 2002, 2: 34-42). Adult neural stem cells are of great potential for repairing damaged or diseased brain tissues, as the germinal centers of the adult brain are free of neuronal damage or dysfunction (Colman A, *Drug Discovery World* 2001, 7: 66-71).

In preferred embodiments, the subject for treatment or prevention, according to the present invention, can be a human, an experimental animal, e.g. a mouse or a rat, a domestic animal, or a non-human primate. The experimental animal can be an animal model for a neurodegenerative disorder, e.g. a transgenic mouse and/or a knock-out mouse with an AD-type neuropathology.

In a further aspect, the invention features a modulator of an activity, or a level, or both said activity and said level of at least one substance which is selected from the group consisting of (i) the TB2 gene, and/or (ii) a transcription product of the TB2 gene and/or (iii) a translation product of the TB2 gene, and/or (iv) a fragment or derivative of (i) to (iii).

In an additional aspect, the invention features a pharmaceutical composition comprising said modulator and preferably a pharmaceutical carrier. Said carrier refers to a diluent, adjuvant, excipient, or vehicle with which the modulator is administered.

In a further aspect, the invention features a modulator of an activity, or a level, or both said activity and said level of at least one substance which is selected from the group consisting of (i) the TB2 gene, and/or (ii) a transcription product of the TB2 gene, and/or (iii) a translation product of the TB2 gene, and/or (iv) a fragment or derivative of (i) to (iii) for use in a pharmaceutical composition.

In another aspect, the invention provides for the use of a modulator of an activity, or a level, or both said activity and said level of at least one substance which is selected from the group consisting of (i) the TB2 gene, and/or (ii) a transcription product of the TB2 gene and/or (iii) a translation product of the TB2 gene, and/or (iv) a fragment or derivative of (i) to (iii) for a preparation of a medicament for treating or preventing a neurodegenerative disease, in particular AD.

In one aspect, the present Invention also provides a kit comprising one or more containers filled with a therapeutically or prophylactically effective amount of said pharmaceutical composition.

In a further aspect, the invention features a recombinant, non-human animal comprising a non-native gene sequence coding for TB2, or a fragment thereof, or a derivative thereof. The generation of said recombinant, non-human animal comprises (i) providing a gene targeting construct containing said gene sequence and a selectable marker sequence, and (ii) introducing said targeting construct into a stem cell of a non-human animal, and (iii) introducing said non-human animal stem cell into a non-human embryo, and (iv) transplanting said embryo into a pseudopregnant non-human animal, and (v) allowing said embryo to develop to term, and (vi) identifying a genetically altered non-human animal whose genome comprises a modification of said gene sequence in both alleles, and (vii) breeding the genetically altered non-human animal of step (vi) to obtain a genetically altered non-human animal whose genome comprises a modification of said endogenous gene, wherein said gene is mis-expressed, or under-expressed, or over-expressed, and

wherein said disruption or alteration results in said non-human animal exhibiting a predisposition to developing symptoms of neuropathology similar to a neurodegenerative disease, in particular AD. Strategies and techniques for the generation and construction of such an animal are known to those of ordinary skill in the art (see e.g. Capecchi, *Science* 1989, 244: 1288-1292 and Hogan et al., 1994, *Manipulating the Mouse Embryo: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York). It is preferred to make use of such a recombinant non-human animal as an animal model for investigating neurodegenerative diseases, in particular AD.

In another aspect, the invention features an assay for screening for a modulator of neurodegenerative diseases, in particular AD, or related diseases and disorders of one or more substances selected from the group consisting of (i) the TB2 gene, and/or (ii) a transcription product of the TB2 gene, and/or (iii) a translation product of the TB2 gene, and/or (iv) a fragment or derivative of (i) to (iii). This screening method comprises (a) contacting a cell with a test compound, and (b) measuring the activity, or the level, or both the activity and the level of one or more substances recited in (i) to (iv), and (c) measuring the activity, or the level, or both the activity and the level of said substances in a control cell not contacted with said test compound, and (d) comparing the levels of the substance in the cells of step (b) and (c), wherein an alteration in the activity and/or level of said substances in the contacted cells indicates that the test compound is a modulator of said diseases and disorders.

In one further aspect, the invention features a screening assay for a modulator of neurodegenerative diseases, in particular AD, or related diseases and disorders of one or more substances selected from the group consisting of (i) the TB2 gene, and/or (ii) a transcription product of the TB2 gene, and/or (iii) a translation product of the TB2 gene, and/or (iv) a fragment or derivative of (i) to (iii), comprising (a) administering a test compound to a test animal which is predisposed to developing or has already developed symptoms of a neurodegenerative disease or related diseases or disorders,

and (b) measuring the activity and/or level of one or more substances recited in (i) to (iv), and (c) measuring the activity and/or level of said substances in a matched control animal which is equally predisposed to developing or has already developed said symptoms and to which animal no such test compound has been administered, and (d) comparing the activity and/or level of the substance in the animals of step (b) and (c), wherein an alteration in the activity and/or level of substances in the test animal indicates that the test compound is a modulator of said diseases and disorders.

In a preferred embodiment, said test animal and/or said control animal is a recombinant, non-human animal which expresses a gene coding for TB2, or a fragment thereof, or a derivative thereof, under the control of a transcriptional regulatory element which is not the native TB2 gene transcriptional control regulatory element.

In another embodiment, the present invention provides a method for producing a medicament comprising the steps of (i) identifying a modulator of neurodegenerative diseases by a method of the aforementioned screening assays and (ii) admixing the modulator with a pharmaceutical carrier. However, said modulator may also be identifiable by other types of screening assays.

In another aspect, the present invention provides for an assay for testing a compound, preferably for screening a plurality of compounds, for inhibition of binding between a ligand and a translation product of the gene coding for TB2, or a fragment or derivative thereof. Said screening assay comprises the steps of (i) adding a liquid suspension of said TB2 translation product, or a fragment or derivative thereof, to a plurality of containers, and (ii) adding a compound or a plurality of compounds to be screened for said inhibition to said plurality of containers, and (iii) adding fluorescently labelled ligand to said containers, and (iv) incubating said TB2 translation product, or said fragment or derivative thereof, and said compound or plurality of compounds, and said fluorescently labelled ligand, and (v) measuring the amounts of

fluorescence associated with said TB2 translation product, or with said fragment or derivative thereof, and (vi) determining the degree of inhibition by one or more of said compounds of binding of said ligand to said TB2 translation product, or said fragment or derivative thereof. It might be preferred to reconstitute said TB2 translation product or fragment or derivative thereof into artificial liposomes to generate the corresponding proteoliposomes to determine the inhibition of binding between a ligand and said TB2 translation product. Methods of reconstitution of membrane proteins from detergent into liposomes are known to those skilled in the art (Banerjee and Datta, *Mol. Cell Biochem.* 1983, 50: 3-15). Instead of utilizing a fluorescently labelled ligand, it might in some aspects be preferred to use any other detectable label known to the person skilled in the art, e.g. radioactive labels, and detect it accordingly. Said method may be useful for the identification of novel compounds as well as for evaluating compounds which have been improved or otherwise optimized in their ability to inhibit the binding of a ligand to a TB2 translation product, or a fragment or derivative thereof.

In one further embodiment, the present invention provides a method for producing a medicament comprising the steps of (i) identifying a compound as an inhibitor of binding between a ligand and a gene product of the gene coding for TB2 by the aforementioned inhibitory binding assay and (ii) admixing the compound with a pharmaceutical carrier. However, said compound may also be identifiable by other types of screening assays.

In another aspect, the invention features an assay for testing a compound, preferably for screening a plurality of compounds to determine the degree of binding of said compounds to a translation product of the gene coding for TB2, or to a fragment or derivative thereof. Said screening assay comprises (i) adding a liquid suspension of said TB2 translation product, or a fragment or derivative thereof, to a plurality of containers, and (ii) adding a fluorescently labelled compound or a plurality of fluorescently labelled compounds to be screened for said binding to said plurality of containers, and

(iii) incubating said TB2 translation product, or said fragment or derivative thereof, and said fluorescently labelled compound or fluorescently labelled compounds, and (iv) measuring the amounts of fluorescence associated with said TB2 translation product, or with said fragment or derivative thereof, and (v) determining the degree of binding by one or more of said compounds to said TB2 translation product, or said fragment or derivative thereof. In this type of assay it might be preferred to use a fluorescent label. However, any other type of detectable label might also be employed. Also in this type of assay it might be preferred to reconstitute a TB2 translation product or fragment or derivative thereof into artificial liposomes as described in the present invention. Said assay methods may be useful for the identification of novel compounds as well as for evaluating compounds which have been improved or otherwise optimized in their ability to bind to a TB2 translation product.

In one further embodiment, the present invention provides a method for producing a medicament comprising the steps of (i) identifying a compound as a binder to a gene product of the TB2 gene by the aforementioned binding assays and (ii) admixing the compound with a pharmaceutical carrier. However, said compound may also be identifiable by other types of screening assays.

In another embodiment, the present invention provides for a medicament obtainable by any of the methods according to the herein claimed screening assays. In one further embodiment, the instant invention provides for a medicament obtained by any of the methods according to the herein claimed screening assays.

The present invention features a protein molecule, said protein molecule being a translation product of the gene coding for TB2, or a fragment, or derivative, or variant thereof, for use as a diagnostic target for detecting a neurodegenerative disease, preferably Alzheimer's disease.

The present invention further features a protein molecule, said protein molecule being a translation product of the gene coding for TB2, or a fragment, or derivative, or variant thereof, for use as a screening target for reagents or compounds preventing, or treating, or ameliorating a neurodegenerative disease, preferably Alzheimer's disease.

The present invention features an antibody which is specifically immunoreactive with an immunogen, wherein said immunogen is a translation product of the TB2 gene, or a fragment, or variant, or derivative thereof. The immunogen may comprise immunogenic or antigenic epitopes or portions of a translation product of said gene, wherein said immunogenic or antigenic portion of a translation product is a polypeptide, and wherein said polypeptide elicits an antibody response in an animal, and wherein said polypeptide is immunospecifically bound by said antibody. Methods for generating antibodies are well known in the art (see Harlow et al., *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York). The term "antibody", as employed in the present invention, encompasses all forms of antibodies known in the art, such as polyclonal, monoclonal, chimeric, recombinatorial, anti-idiotypic, humanized, or single chain antibodies, as well as fragments thereof. Antibodies of the present invention are useful, for instance, in a variety of diagnostic and therapeutic methods involving detecting translation products of the TB2 gene.

In a preferred embodiment of the present invention, said antibodies can be used for detecting the pathological state of a cell in a sample from a subject, comprising immunocytochemical staining of said cell with said antibody, wherein an altered degree of staining, or an altered staining pattern in said cell compared to a cell representing a known health status indicates a pathological state of said cell. Preferably, the pathological state relates to a neurodegenerative disease, in particular to AD. Immunocytochemical staining of a cell can be carried out by a number of different experimental methods well known in the art. It might be preferred, however, to apply an automated method for the detection of antibody binding, wherein the determination of the

degree of staining of a cell, or the determination of the cellular or subcellular staining pattern of a cell, or the topological distribution of an antigen on the cell surface or among organelles and other subcellular structures within the cell, are carried out according to the method described in US patent 6150173.

Other features and advantages of the invention will be apparent from the following description of figures and examples which are illustrative only and not intended to limit the remainder of the disclosure in any way.

Figure 1 depicts the brain regions with selective vulnerability to neuronal loss and degeneration in AD. Primarily, neurons within the inferior temporal lobe, the entorhinal cortex, the hippocampus, and the amygdala are subject to degenerative processes in AD (Terry et al., *Annals of Neurology* 1981, 10:184-192). These brain regions are mostly involved in the processing of learning and memory functions. In contrast, neurons within the frontal cortex, the occipital cortex, and the cerebellum remain largely intact and preserved from neurodegenerative processes in AD. Brain tissues from the frontal cortex (F) and the temporal cortex (T) of AD patients and healthy, age-matched control individuals were used for the herein disclosed examples. For illustrative purposes, the image of a normal healthy brain was taken from a publication by Strange (*Brain Biochemistry and Brain Disorders*, Oxford University Press, Oxford, 1992, p.4).

Figure 2 illustrates the verification of the differential expression of the TB2 gene in AD brain tissues by quantitative RT-PCR analysis. Quantification of RT-PCR products from RNA samples collected from the frontal cortex (F) and temporal cortex (T) of an AD patient (Fig 2A) and of a healthy, age-matched control individuals (Fig 2B) was performed by the LightCycler rapid thermal cycling technique. The data were normalized to the combined average values of a set of standard genes which showed no significant differences in their gene expression levels. Said set of standard genes consisted of genes for the ribosomal protein S9, the transferrin receptor, GAPDH, and beta-actin. The

figure depicts the kinetics of amplification by plotting the cycle number against the amount of amplified material as measured by its fluorescence. Note that the amplification kinetics of the TB2 cDNA from both the frontal and temporal cortices of a normal control individual during the exponential phase of the reaction overlap (Fig 2B, arrow), whereas in AD (Fig 2A, arrows), there is a significant separation of the corresponding curves, indicating a differential expression of the TB2 gene in the two analyzed brain regions.

Figure 3 discloses SEQ ID NO.1, the nucleotide sequence of the 461 bp TB2 cDNA fragment, identified and obtained by subtractive suppressive microarray hybridization experiments and subsequent cloning.

Figure 4 charts the schematic alignment of SEQ ID No.1 with the nucleotide sequence of human TB2 mRNA/cDNA (Kinzler et al, *Science* 1991, 253: 661-665; GenBank accession number BC000232); polyA: polyadenylation site; the open box represents the TB2 open reading frame (positions 38-595).

Figure 5 displays the exact sequence alignment of SEQ ID NO.1 with the nucleotide sequence of the human TB2 mRNA/cDNA (Kinzler et al, *Science* 1991, 253: 661-665; GenBank accession number BC000232).

Table 1 discloses the initial identification of differential expression of the human TB2 gene by subtractive suppressive microarray hybridization experiments. Identical biochips containing cDNA clones of subtracted AD and control brain cDNA libraries were co-hybridized with different Cyanine3 (Cy3) and Cyanine5 (Cy5) labeled SMART cDNA probes derived from frontal or temporal cortex of AD patients and control persons, respectively. For details of probe production, see section (vi). The table lists the gene expression level of TB2 as the ratio of fluorescence intensity measured for the temporal cortex of control persons relative to the temporal cortex of AD patients (biochip 5) and the frontal cortex of control persons relative to the frontal cortex of AD patients (biochip 7). The ratios of fluorescence intensity reflect an up-regulation of TB2 gene expression in the frontal cortex or a down-regulation

of TB2 gene expression in the temporal cortex of AD patients in comparison to non-AD healthy control individuals.

Table 2 lists the gene expression levels in the frontal cortex relative to the temporal cortex for the TB2 gene in seven AD patients (0.59 to 1.99 fold) and five healthy, age-matched control individuals (0.98 to 1.26 fold). The values shown are reciprocal values according to the formula described herein (see below).

EXAMPLE I:

(i) Brain tissue dissection from patients with AD:

Brain tissues from AD patients and age-matched control subjects were collected within 6 hours post-mortem and immediately frozen on dry ice. Sample sections from each tissue were fixed in paraformaldehyde for histopathological confirmation of the diagnosis. Brain areas for differential expression analysis were identified (see Fig. 1) and stored at -80°C until RNA extractions were performed.

(ii) Isolation of total mRNA:

Total RNA was extracted from post-mortem brain tissue by using the RNeasy kit (Qiagen) according to the manufacturer's protocol. The accurate RNA concentration and the RNA quality were determined with the DNA LabChip system using the Agilent 2100 Bioanalyzer (Agilent Technologies). For additional quality testing of the prepared RNA, i.e. exclusion of partial degradation and testing for DNA contamination, specifically designed intronic GAPDH oligonucleotides and genomic DNA as reference control were used to generate a melting curve with the LightCycler technology as described in the manufacturer's protocol (Roche).

(iii) cDNA synthesis and Rsa I digestion:

In order to identify changes in gene expression in different tissues, a screening method combining cDNA synthesis, suppressive subtractive hybridization (SSH) and screening of microarray chips with a diversity of cDNA probes from SSH was employed. This technique compares different populations of mRNA and provides clones of genes that are expressed in one population of cells but not, or at lower level, in the other population of cells. In the present invention, RNA populations from selected post-mortem brain tissues (frontal and temporal cortex) of AD patients and age-matched control subjects were compared.

As starting material for the suppressive subtractive microarray analysis total RNA was extracted as described above (ii). For production of preferably full-length cDNAs, the polymerase chain reaction (PCR)-based method 'SMART cDNA Synthesis' was performed according to the manufacturer's protocol (Clontech).

The principle of 'SMART cDNA synthesis' has been described in detail (Chenchik et al., in *Gene Cloning and Analysis by RT-PCR*, Eds. Siebert and Larrick, *Biotechniques Books, Natick, MA* 1998: 305-320). For SMART cDNA synthesis, four RNA pools, each consisting of 8 µg total RNA, were prepared. Each pool contained 2 µg of each of four different samples, i.e. from inferior frontal cortex (CF) and from inferior temporal cortex (CT) of control brains, from inferior frontal cortex (PF) and from inferior temporal cortex (PT) of patient brains, respectively. An amount of 1 µg of total RNA mix was utilized in a reaction volume of 50 µl (PCR cycler: Multi Cyclor PTC 200, MJ Research). The second SMART PCR step was performed using 19 cycles. SuperScript II RNaseH Reverse Transcriptase and 5x first-strand buffer (Invitrogen) were used.

After extraction and purification of the PCR products, restriction digestions were carried out with 30 U Rsa I (MBI Fermentas) for 2.5 hours at 37°C. Rsa I restriction sites are located within the universal priming sites of the double stranded (ds) cDNA. The quality of the digestions was analyzed by agarose gel electrophoresis, the digested samples were purified (QIAquick PCR Purification Kit, Qiagen), and the cDNA concentrations were determined by UV spectrophotometry (Biorad).

(iv) suppressive subtractive hybridization (SSH):

Four SMART cDNA pools (iii), were compared using suppressive subtractive hybridization. A pool of cDNA containing differentially expressed genes is thereby designated as "Tester", the reference cDNA pool as "Driver". The two pools are hybridized, and all cDNAs present in both pools, will be eliminated, i.e. the Driver-pool will be subtracted from the Tester-pool. Thus, clones of genes that are predominantly expressed in the Tester population are obtained.

The 'PCR-Select cDNA Subtraction Kit' was used to perform the subtractive hybridization (Clontech). The 'Tester' SMART cDNA pools, derived from temporal cortex (CT) of control brains, and from frontal and temporal cortex (PF and PT) of patient brains (III), were subdivided into two pools each. Each pool was ligated with Adaptor 1 or Adaptor 2, respectively, thus obtaining 6 different 'Tester' cDNA pools. The three 'Driver' SMART cDNA pools, CT, PF and PT, remained unligated. In a first hybridization step, used to enrich for differentially expressed sequences, the following three different 'Tester' SMART cDNA pools were combined with an excess of the following 'Driver' SMART cDNAs: SSH(1): PF-'Tester' and PT-'Driver'; SSH(2): PT-'Tester' and PF-'Driver'; SSH(3): CT-'Tester' and PT-'Driver'; SSH(4): PT-'Tester' and CT-'Driver'. Following a denaturation step for 1.5 min at 98°C, the hybridization was carried out for 8 hours at 68°C. In a second step, the two corresponding primary hybridization samples of 'Tester' SMART cDNA pools ligated to Adaptor 1 or 2, respectively, were mixed and re-hybridized at 68°C for 15 hours, with an excess of the 'Driver' SMART cDNA pool, as used before. Thus, suitable double stranded cDNAs for subsequent amplification, i.e. with both adaptor sequences at their 5' and 3' ends and therefore with different annealing sites, were generated. The following PCR steps were applied to obtain efficiently amplified specific products and to suppress nonspecific amplification. In the first PCR, missing strands of the adaptors were filled in by DNA-polymerase activity. 1 µl of the obtained hybridization products each were subjected to PCR using the corresponding 'Primer 1' (10 µM) (Clontech) along with 1x PCR reaction buffer (Clontech), 10 mM dNTP-Mix (dATP, dGTP,

dCTP, dTTP, Amersham Pharmacia Biotech), and 0.5 µl 50x Advantage cDNA Polymerase Mix (Clontech) in a 25 µl final volume. PCR conditions were set as follows: one round at 75 °C for 5 min, which was followed by 27 or 30 cycles: 94 °C for 30 sec, 64 °C or 66 °C for 30 sec, 72 °C for 1.5 min. One final step at 72 °C for 5 min was added to the last cycle. A second nested PCR was performed as described for the first PCR, except that instead of 'Primer 1' the nested primers 'Nested Primer 1' and '2R' were used and an annealing temperature of 66°C or 68°C and 12 or 15 cycles were applied. PCR-products obtained by different conditions were pooled for subsequent analysis. For the primer sequences used, refer to appendix B of the supplier's user manual (Clontech).

(v) Cloning of subtracted PCR products and production of DNA-biochips:
The SSH SMART double stranded cDNAs of the four different combinations SSH(1)-SSH(4), refer to (iv), were ligated into the pCR2.1-vector and transformed into INValphaF' cells according to the manufacturer's instructions (TA Cloning Kit, Invitrogen). Bacterial colonies were picked and analyzed by colony PCR on MTPs (microtiter plates, 96 well, Abgene), using 'Nested Primer 1' and 'Nested Primer 2'. Those MTPs showing more than 90% positive clones were subjected to a preparative colony PCR approach. Per well, the following PCR mix was generated: the corresponding oligonucleotides 'Nested Primer 1' and 'Nested Primer 2' (0.5 µM each), 1 x Titanium PCR buffer (Clontech), 200 µM dNTP-Mix (Amersham Pharmacia Biotech), 0.2 x TitaniumTaq DNA-Polymerase (Clontech) in a 120 µl final volume. PCR conditions were set as follows: one round at 94 °C for 30 sec for denaturing, the next round was followed by 35 cycles: 94 °C for 30 sec and 68°C for 3 min. The quality of the amplified products was checked and analyzed (DNA LabChip system, Agilent 2100 Bioanalyzer, Agilent Technologies), followed by purification (Multiscreen-PCR-Purification system, Millipore).

Additionally, the following standard control samples were generated: three different *Arabidopsis thaliana* genes, polyA-DNA, salmon sperm DNA, human Cot-1 DNA, and 3xSSC-buffer were used as negative controls (Microarray

Validation System, Stratagene); beta-Actin and Xenopus cDNA were used as normalizing controls.

Several MTPs were made of each of the SSH combinations SSH(1)-(4), harboring amplification products of 96 different clones per plate. The amplified products were spotted in triplicates onto GAPS glass-slides (CMT-GAPS, Corning) by GeneScan Europe.

(vi) Probe synthesis and identification of differentially expressed genes by screening of DNA biochips:

For the production of Cyanine3 (Cy3) and Cyanine5 (Cy5) labeled SMART cDNA-probes, the PCR-based method 'SMART cDNA Synthesis' was performed as described in section (iii). Here, we used total RNA as starting material, which was extracted as described above (ii). Four RNA mixtures were prepared as described in section (iii). 1 µg of each RNA mix and 1 ng Xenopus total RNA were subjected to the SMART cDNA reaction. For PCR amplification, extraction and purification of the cDNAs, restriction digestion with Rsa I, and subsequent purification of the digested samples, refer to section (iii).

SMART cDNA samples were labeled with either Cy3 or Cy5 (Atlas Glass Fluorescent Labeling Kit, Clontech). In the first labeling step, aliphatic amino groups, i.e. aminoallyl-dUTP (Clontech), were incorporated into denatured (100°C, 7 min) Rsa I digested PCR products. The reaction was catalyzed by the Klenow Fragment (MBI Fermentas). In a second labeling step, the fluorescent reporter dyes Cy3 or Cy5 were coupled to the incorporated functionalities.

The purified Cy3 and Cy5 labeled SMART cDNA probes (Atlas NucleoSpin Extraction Kit, Clontech) were resuspended in hybridization buffer (5x SSC, 0.1% SDS, 25% formamide) after denaturation for 7 min at 100°C. Subsequently, the Cy3 labeled SMART probe was mixed with the Cy5 labeled SMART probe and together applied evenly onto one prehybridized (5x SSC, 0.1% SDS, 1% BSA, 45 min at 42°C) microarray. Array hybridization was performed over night at 42°C. High stringency washing of the biochips

followed according to the instructions of the TSA Detection Kit protocol (NEN Life Science).

Biochip 5 was hybridized with SMART cDNA mix PT (Cy3) and CT (Cy5), biochip 7 was hybridized with SMART cDNA mix PF (Cy3) and CF (Cy5), respectively. Scanning the microarrays with the appropriate wavelengths (635 nm, 532 nm) allowed detection of both cyanine dyes simultaneously.

(vii) DNA biochips data evaluation:

Fluorescence raw data for Cy3 and Cy5, measured at 635 and 532 nm, respectively, were taken severalfold (for each of the three spots per cDNA). One set of measurements was performed within the spot area (signal) and another set of measurements was taken nearby (background). Subsequently the net fluorescence intensity (FI_{635} , FI_{532}) of the spots was calculated as follows:

$$FI_{635/532} = (M FI_{spot} - 1 SD FI_{spot}) - (M FI_{background} + 1 SD FI_{background}).$$

In this calculation, M defines the median of the replicate measurements per spot, SD the standard deviation of the corresponding mean. Subsequently, only FI_{635} and FI_{532} values of >2 were considered for further evaluation plus those FI_{635} and FI_{532} values of <2 where the corresponding value for the second wavelength was >3 .

In an analogous manner, the corresponding values for the *Xenopus* cDNA control and the set of standard (housekeeping) genes were evaluated. The *Xenopus* cDNA was used as an internal calibrator for the efficiency of cDNA synthesis of the disease relevant mRNAs. Then, from the background corrected $FI_{635/532}$ medians of the three replicate spots, the statistical mean was calculated and the signal ratio R for the cDNA probes was derived using formula :

$$R_{635/532} = FI_{635, calibrated} / FI_{532, calibrated}.$$

In a last step of evaluation, the results of the different hybridizations were considered for logical coherence.

(viii) Confirmation of differential expression by quantitative RT-PCR:

The expression levels of the human TB2 gene in frontal versus temporal cortex were analyzed using the LightCycler technology (Roche). This technique features rapid thermal cycling for the polymerase chain reaction as well as real-time measurement of fluorescent signals during amplification and therefore allows for highly accurate quantification of RT-PCR products by using a kinetic, rather than an endpoint readout. The ratio of TB2 cDNA in frontal cortex versus temporal cortex was determined (relative quantification).

First, a standard curve was generated to determine the efficiency of the PCR with specific primers for human TB2 (5'-CGAGTCCCAGATGGACAGTGT-3' and 5'-GTGATGGCATCTGCAGTCTCTT-3'). PCR amplification (95 °C and 1 sec, 56 °C and 5 sec, and 72 °C and 5 sec) was performed in a volume of 20 µl containing LightCycler-FastStart DNA Master SYBR Green I mix (FastStart Taq DNA polymerase, reaction buffer, dNTP mix with dUTP instead of dTTP, SYBR Green I dye, and 1 mM MgCl₂; Roche), 0.5 µM primers, 2 µl of a cDNA dilution series (final concentration of 40, 20, 10, 5, 1 and 0.5 ng human total brain cDNA; Clontech) and, depending on the primers used, additional 3 mM MgCl₂. Melting curve analysis revealed a single peak at approximately 82.5°C with no visible primer dimers. Quality and size of the PCR product were determined with the DNA LabChip system (Agilent 2100 Bioanalyzer, Agilent Technologies). A single peak at the expected size of 69 bp for TB2 was observed in the electropherogram of the sample.

In an analogous manner, the PCR protocol was applied to determine the PCR efficiency of a set of reference genes which were selected as a reference standard for quantification. In the present invention, the mean value of five such reference genes was determined: (1) cyclophilin B, using the specific primers 5'-ACTGAAGCACTACGGGCCTG-3' and 5'-AGCCGTTGGTGTCTT-TGCC-3' except for MgCl₂ (an additional 1 mM was added instead of 3 mM). Melting curve analysis revealed a single peak at approximately 87 °C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band of the expected size (62 bp). (2) Ribosomal protein S9 (RPS9),

using the specific primers 5'-GGTCAAATTTACCCTGGCCA-3' and 5'-TCTCATCAAGCGTCAGCAGTTC-3' (exception: additional 1 mM MgCl₂ was added instead of 3 mM). Melting curve analysis revealed a single peak at approximately 85°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (62 bp). (3) beta-actin, using the specific primers 5'-TGGAACGGTGAAGGTGACA-3' and 5'-GGCAAGGGACTTCCTGTAA-3'. Melting curve analysis revealed a single peak at approximately 87°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (142 bp). (4) GAPDH, using the specific primers 5'-CGTCATGGGTGTGAACCATG-3' and 5'-GCTAAGCAGTTGGTGGTGACAG-3'. Melting curve analysis revealed a single peak at approximately 83°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (81 bp). (5) Transferrin receptor TRR, using the specific primers 5'-GTCGCTGGTCAGTTCGTGATT-3' and 5'-AGCAGTTGGCTGTTGTACCTCTC-3'. Melting curve analysis revealed a single peak at approximately 83°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (80 bp).

For calculation of the values, first the logarithm of the cDNA concentration was plotted against the threshold cycle number C_t for TB2 and the five reference standard genes. The slopes and the intercepts of the standard curves (i.e. linear regressions) were calculated for all genes. In a second step, cDNAs from frontal cortex and temporal cortex were analyzed in parallel and normalized to cyclophilin B. The C_t values were measured and converted to ng total brain cDNA using the corresponding standard curves:

$$10^{((C_t \text{ value} - \text{intercept}) / \text{slope})} \quad [\text{ng total brain cDNA}]$$

The values for frontal and temporal cortex cDNAs of TB2 cDNA were normalized to cyclophilin B and the ratio was calculated according to formula:

$$\text{Ratio} = \frac{\text{TB2 temporal [ng] / cyclophilin B temporal [ng]}}{\text{TB2 frontal [ng] / cyclophilin B frontal [ng]}}$$

In a third step, the set of reference standard genes was analyzed in parallel to determine the mean average value of the temporal to frontal ratios of expression levels of the reference standard genes for each individual brain sample. As cyclophilin B was analyzed in step 2 and step 3, and the ratio from one gene to another gene remained constant in different runs, it was possible to normalize the values for TB2 to the mean average value of the set of reference standard genes instead of normalizing to one single gene alone. The calculation was performed by dividing the ratio shown above by the deviation of cyclophilin B from the mean value of all housekeeping genes. The results of one such quantitative RT-PCR analysis for the TB2 gene are shown in Figure 2.

CLAIMS

1. A method of diagnosing or prognosticating a neurodegenerative disease in a subject, or determining whether a subject is at increased risk of developing said disease, comprising:

determining a level and/or an activity of

- (i) a transcription product of the TB2 gene, and/or
 - (ii) a translation product of the TB2 gene, and/or
 - (iii) a fragment or derivative of said transcription or translation product
- in a sample from said subject and comparing said level and/or said activity to a reference value representing a known disease or health status, thereby diagnosing or prognosticating said neurodegenerative disease in said subject, or determining whether said subject is at increased risk of developing said neurodegenerative disease.

2. The method according to claim 1 wherein said neurodegenerative disease is Alzheimer's disease.

3. A kit for diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, in a subject, or determining the propensity or predisposition of a subject to develop such a disease, said kit comprising:

- (a) at least one reagent which is selected from the group consisting of (i) reagents that selectively detect a transcription product of the TB2 gene and (ii) reagents that selectively detect a translation product of the TB2 gene; and
- (b) an instruction for diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, or determining the propensity or predisposition of a subject to develop such a disease by (i) detecting a level, or an activity, or both said level and said activity, of said transcription product and/or said translation product of the TB2 gene, in a sample from said subject; and (ii) diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, or determining the propensity or predisposition of said subject to develop

such a disease, wherein a varied level, or activity, or both said level and said activity, of said transcription product and/or said translation product compared to a reference value representing a known health status; or a level, or activity, or both said level and said activity, of said transcription product and/or said translation product similar or equal to a reference value representing a known disease status indicates a diagnosis or prognosis of a neurodegenerative disease, in particular Alzheimer's disease, or an increased propensity or predisposition of developing such a disease.

4. A modulator of an activity and/or of a level of at least one substance which is selected from the group consisting of:

- (i) the TB2 gene, and/or
- (ii) a transcription product of the TB2 gene, and/or
- (iii) a translation product of the TB2 gene, and/or
- (iv) a fragment or derivative of (i) to (iii)

for use in a pharmaceutical composition.

5. A recombinant, non-human animal comprising a non-native gene sequence coding for TB2 or a fragment, or a derivative, or a variant thereof, said animal being obtainable by:

- (i) providing a gene targeting construct comprising said gene sequence and a selectable marker sequence, and
- (ii) introducing said targeting construct into a stem cell of a non-human animal, and
- (iii) introducing said non-human animal stem cell into a non-human embryo, and
- (iv) transplanting said embryo into a pseudopregnant non-human animal, and
- (v) allowing said embryo to develop to term, and
- (vi) identifying a genetically altered non-human animal whose genome comprises a modification of said gene sequence in both alleles, and

- (vii) breeding the genetically altered non-human animal of step (vi) to obtain a genetically altered non-human animal whose genome comprises a modification of said endogenous gene, wherein said disruption results in said non-human animal exhibiting a predisposition to developing symptoms of a neurodegenerative disease or related diseases or disorders.

6. An assay for screening for a modulator of neurodegenerative diseases, in particular Alzheimer's disease, or related diseases or disorders of one or more substances selected from the group consisting of:

- (i) the TB2 gene, and/or
- (ii) a transcription product of the TB2 gene, and/or
- (iii) a translation product of the TB2 gene, and/or
- (iv) a fragment or derivative of (i) to (iii),

said method comprising:

- (a) contacting a cell with a test compound;
- (b) measuring the activity and/or level of one or more substances recited in (i) to (iv);
- (c) measuring the activity and/or level of one or more substances recited in (i) to (iv) in a control cell not contacted with said test compound; and
- (d) comparing the levels and/or activities of the substance in the cells of step (b) and (c), wherein an alteration in the activity and/or level of substances in the contacted cells indicates that the test compound is a modulator of said diseases or disorders.

7. A method of screening for a modulator of neurodegenerative diseases, in particular Alzheimer's disease, or related diseases or disorders of one or more substances selected from the group consisting of

- (i) the TB2 gene, and/or
- (ii) a transcription product of the TB2 gene, and/or
- (iii) a translation product of the TB2 gene, and/or
- (v) a fragment or derivative of (i) to (iii),

said method comprising:

- (a) administering a test compound to a test animal which is predisposed to developing or has already developed symptoms of a neurodegenerative disease or related diseases or disorders in respect of the substances recited in (i) to (iv);
- (b) measuring the activity and/or level of one or more substances recited in (i) to (iv);
- (c) measuring the activity and/or level of one or more substances recited in (i) or (iv) in a matched control animal which is predisposed to developing or has already developed symptoms of a neurodegenerative disease or related diseases or disorders in respect to the substances recited in (i) to (iv) and to which animal no such test compound has been administered;
- (d) comparing the activity and/or level of the substance in the animals of step (b) and (c), wherein an alteration in the activity and/or level of substances in the test animal indicates that the test compound is a modulator of said diseases or disorders.

8. The method according to claim 7 wherein said test animal and/or said control animal is a recombinant animal which expresses a gene coding for TB2, or a fragment thereof, or a derivative thereof, under the control of a transcriptional control element which is not the native TB2 gene transcriptional control element.

9. An assay for testing a compound, preferably for screening a plurality of compounds for inhibition of binding between a ligand and a translation product of the gene coding for TB2, or a fragment or derivative thereof, said assay comprising the steps of:

- (i) adding a liquid suspension of said TB2 translation product, or a fragment or derivative thereof, to a plurality of containers;
- (ii) adding a compound or a plurality of compounds to be screened for said inhibition to said plurality of containers;
- (iii) adding fluorescently labelled ligand to said containers;

- (iv) incubating said TB2 translation product, or said fragment or derivative thereof, and said compound or compounds, and said fluorescently labelled ligand;
- (v) measuring amounts of fluorescence associated with said TB2 translation product, or with said fragment or derivative thereof; and
- (vi) determining the degree of inhibition by one or more of said compounds of binding of said ligand to said TB2 translation product, or said fragment or derivative thereof.

10. A protein molecule, said protein molecule being a translation product of the gene coding for TB2, or a fragment, or derivative, or variant thereof, for use as a diagnostic target for detecting a neurodegenerative disease, preferably Alzheimer's disease.

11. A protein molecule, said protein molecule being a translation product of the gene coding for TB2, or a fragment, or derivative, or variant thereof, for use as a screening target for reagents or compounds preventing, or treating, or ameliorating a neurodegenerative disease, preferably Alzheimer's disease.


12. Use of an antibody specifically immunoreactive with an immunogen, wherein said immunogen is a translation product of the gene coding for TB2, or a fragment, or derivative, or variant thereof, for detecting a pathological state of a cell in a sample from a subject, comprising immunocytochemical staining of said cell with said antibody, wherein an altered degree of staining, or an altered staining pattern in said cell compared to a cell representing a known health status indicates a pathological state of said cell.

SUMMARY

The present invention discloses the differential expression of the TB2 gene in specific brain regions of Alzheimer's disease patients. Based on this finding, this invention provides a method for diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, in a subject, or for determining whether a subject is at increased risk of developing such a disease. Furthermore, this invention provides therapeutic and prophylactic methods for treating or preventing Alzheimer's disease and related neurodegenerative disorders using the TB2 gene and its corresponding gene products. A method of screening for modulating agents of neurodegenerative diseases is also disclosed.


- 1 / 7 -

Figure 1: Identification of genes involved in Alzheimer's Disease pathology



- 2 / 7 -

Figure 2: Differential expression of the TB2 gene as determined by RT-PCR analysis




- 3 / 7 -

Figure 3: SEQ ID NO.1

```
1  ACCTGGTGTT CGGTTATGGG GCCTCTCTCC TCTGCAACCT GATAGGATTT
51  GGCTACCCAG CCTACATCTC AATTAAAGCT ATAGAGAGTC CCAACAAAGA
101 AGATGATACC CAGTGGCTGA CCTACTGGGT AGTGTATGGT GTGTTCAAGCA
151 TTGCTGAATT CTTCTCTGAT ATCTTCCTGT CATGGTTCCC CTTCTACTAC
201 ATGCTGAAGT GTGGCTTCCT GTTGTGGTGC ATGGCCCCGA GCCTTCTAAT
251 GGGGCTGAAC TGCTCTACAA GCGCATCATC CGGCCTTTCT TCCTGAAGCA
301 CGAGTCCCAG ATGGACAGTG TGGTCAAGGA CCTTAAAGAC AAGGCCAAAG
351 AGACTGCAGA TGCCATCACT AAAGAAGCGA AGAAAGCTAC CGTGAATTTA
401 CTGGGTGAAG AAAAGAAGAG CACCTAAACC AGACTGGATG GAAACTTCCT
451 GCCCTCTCTG T
```

- 4 / 7 -

**Figure 4: Schematic alignment of SEQ ID NO.1 to human TB2
cDNA (GenBank accession number BC000232)**



- 5 / 7 -

**Figure 5: Sequence alignment of SEQ ID NO.1 to
nucleotides 168-629 of human TB2 cDNA
(GenBank accession number BC000232)**

```
1  ACCTGGTGTTCGGTTATGGGGCCTCTCTCCTCTGCAACCTGATAGGATTT 50
   ||||||||||||||||||| |||||||||||||||||||
168 ACCTGGTGTTCGGTTATGGAGCCTCTCTCCTCTGCAACCTGATAGGATTT 217
   ||||||||||||||||||| |||||||||||||||||||
51  GGCTACCCAGCCTACATCTCAATTAAAGCTATAGAGAGTCCCAACAAAGA 100
   ||||||||||||||||||| |||||||||||||||||||
218 GGCTACCCAGCCTACATCTCAATTAAAGCTATAGAGAGTCCCAACAAAGA 267
   ||||||||||||||||||| |||||||||||||||||||
101 AGATGATACCCAGTGGCTGACCTACTGGGTAGTGTATGGTGTGTTTCAGCA 150
   ||||||||||||||||||| |||||||||||||||||||
268 AGATGATACCCAGTGGCTGACCTACTGGGTAGTGTATGGTGTGTTTCAGCA 317
   ||||||||||||||||||| |||||||||||||||||||
151 TTGCTGAATTCTTCTCTGATATCTTCCTGTGTCATGGTTCCCCTTCTACTAC 200
   ||||||||||||||||||| |||||||||||||||||||
318 TTGCTGAATTCTTCTCTGATATCTTCCTGTGTCATGGTTCCCCTTCTACTAC 367
   ||||||||||||||||||| |||||||||||||||||||
201 ATGCTGAAGTGTGGCTTCCTGTTGTGGTGCATGGCCCCGAG.CCTTCTAA 249
   ||||||||||||||||||| |||||||||||||||||||
368 ATGCTGAAGTGTGGCTTCCTGTTGTGGTGCATGGCCCCGAGCCCTTCTAA 417
   ||||||||||||||||||| |||||||||||||||||||
250 TGGGGCTGAACTGCTCTACAAGCGCATCATCCGGCCTTTCTTCCTGAAGC 299
   ||||||||||||||||||| |||||||||||||||||||
418 TGGGGCTGAACTGCTCTACAAGCGCATCATCCGTCCTTTCTTCCTGAAGC 467
   ||||||||||||||||||| |||||||||||||||||||
300 ACGAGTCCCAGATGGACAGTGTGGTCAAGGACCTTAAAGACAAGGCCAAA 349
   ||||||||||||||||||| |||||||||||||||||||
468 ACGAGTCCCAGATGGACAGTGTGGTCAAGGACCTTAAAGACAAGGCCAAA 517
   ||||||||||||||||||| |||||||||||||||||||
350 GAGACTGCAGATGCCATCACTAAAGAAGCGAAGAAAGCTACCGTGAATTT 399
   ||||||||||||||||||| |||||||||||||||||||
518 GAGACTGCAGATGCCATCACTAAAGAAGCGAAGAAAGCTACCGTGAATTT 567
   ||||||||||||||||||| |||||||||||||||||||
400 ACTGGGTGAAGAAAAGAAGAGCACCTAAACCAGACTGGATGGAAACTTCC 449
   ||||||||||||||||||| |||||||||||||||||||
568 ACTGGGTGAAGAAAAGAAGAGCACCTAAACCAGACTGGATGGAAACTTCC 617
   ||||||||||||||||||| |||||||||||||||||||
450 TGCCCTCTCTGT 461
   |||||||||||
618 TGCCCTCTCTGT 629
```

- 6 / 7 -

Table 1: Identification of differentially expressed genes in microarray hybridization experiments

Biochip	Type of probe	Used probes (Cy5-/Cy3- labeled)	Ratio fluorescence intensity:
5	SMART cDNA	CT / PT	2.91
7	SMART cDNA	CF / PF	0.49

- 7 / 7 -

Table 2: Differential expression of the TB2 gene in the Alzheimer's disease brain

sample	Δfold (frontal / temporal cortex)
control 1	1.26
control 2	1.09
control 3	1.00
control 4	0.99
control 5	0.98
patient 1	1.99
patient 2	1.60
patient 3	0.59
patient 4	1.38
patient 5	0.95
patient 6	1.57
patient 7	1.78

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.